



RAINFALL INTERCEPTION FOR SIXTY-YEAR-OLD PINE STAND AT THE TUCZNO FOREST DISTRICT

*Marek Urbaniak, Janusz Olejnik, Antoni T. Miler,
Anna Krysztofiak-Kaniewska, Klaudia Ziemblińska*
Poznań University of Life Sciences

Summary

The aim of the presented studies was to calculate interception values for the sixty-year-old pine stand located in Tuczno Forest District. The area subjected to analysis is located in the north-western part of Poland, in West Pomerania province. The calculations were based on the measurements of precipitation above the tree crowns (2 pluviometers) and at a height of 1 m above the surface (13 pluviometers). Precipitation measurements used in this work were carried out in 30-minute intervals during the entire 2014. The analyses were conducted using mainly A-STER tipping bucket pluviometers. Temporary interception in the studied sixty-year-old pine stand can reach even about seven millimeters. The daily difference between precipitation above the tree crowns and precipitation at 1 m above the surface can reach twelve millimeters. The calculated volume of the so called interception reservoir was about three millimeters (3.26 mm). The annual value of interception for the tested stand was 19.6% of the total annual precipitation.

Key words: interception, forest meteorology, monitoring-gauging station Tuczno

INTRODUCTION

In field conditions, forest rainfall interception is defined as a difference between the precipitation measured above the tree crowns and the precipitation measured under tree cover (Calder 2001, Bryant et al. 2005, Pypker et al. 2005).

Attempts at mathematical description of interception process have been made for about fifty years (Czarnowski, Olszewski 1968, Crocford and Richardson 2000, Keim et al. 2004). They concerned particularly the interception for individual forest zones (Jong and Jetten 2007), the forest canopy (Gash et al 1995), or a component of water balance (Pike and Scherer 2003).

Studies on mathematical model of rainfall interception, both based on physical relationships (Rutter and Morton 1977) and conceptual ones (Suliński 1993) base on empirical laboratory or field measurements. The conceptual models usually assume the interception by trees may be identified with a process of “filling a leaky reservoir” (Osuch et al. 2005, Toba and Ohta 2008). The main affecting factors are: sprayed area (LAI index) (Suliński 1993, Jong and Jetten 2007) and rainfall characteristics (the rainfall intensity and the size of falling drops) (Keim et al. 2006).

A review description of various measurement methods applied so far and conceptions of mathematical descriptions of interception by trees and forest vegetation may be found e.g. in the paper by Klamerus-Iwan (2014).

This paper aims at presentation of a short lived (associated with precipitation episodes) and seasonal interception for sixty-year-old pine stand located in the central, lowland part of Poland.

METHODS

A measurement tower, 34-m high, was constructed in the centre of TUCZNO research facility of the Department of Meteorology, Poznań University of Life Sciences. Two pluviometers were installed on its top, one A-STER tipping basket and the other forming a part of Vaisala meteorological station. Thirteen A-STER tipping basket pluviometers were arranged on a cross plan (N-S, E-W) in the forest bottom, on a standard height of 1m above the surface. Precipitation was summed up in 30 minute intervals for all 15 pluviometers (Olejnik et al. 2013, Urbaniak et al. 2014).

DESCRIPTION OF THE OBJECT OF RESEARCH

The described experimental area is located in the north-western part of Poland in Tuczno Forest District, in the area of Martew forest inspectorate. The terrain is situated in the Drawa National Park buffer zone. Formerly it was an agricultural land, later afforested and now over sixty-year-old pine stand is growing on mixed forest habitat. The share of pine (*Pinus sylvestris* L.) in species composition of the stand reaches almost 99%. The remaining 1% is an admixture of silver birch (*Betula pendula* Roth.). The underbrush is composed mainly of beech (*Fagus sylvatica* L.) and hornbeam (*Carpinus betulus* L.). The soils were

identified as proper rusty soil, and rusty gley soil in some places, deposited on loamy sand (Chojnicki et al. 2009).

RESULTS

Precipitations measured in 30 minute intervals above the tree crowns (PG) and at the height of 1 m above the surface (PD) were shown in Figure 1. These values are means for the measurements from respectively two and 13 pluviometers. The maximum value of PG was 13.70 mm ($\sigma_{(PG)} = 0.26$ mm), whereas the maximum value of PD was 6.53 mm ($\sigma_{(PD)} = 0.16$ mm). This proportion results from a high temporary interception value.

The daily PG and PD precipitation totals were shown in Figure 2. The maximum difference between PG and PD daily totals is 11.96 mm ($\sigma_{(PG-PD)} = 1.61$ mm).




Figure 1. Precipitation measured at 1 m above the surface (red line) vs. measured above the forest canopy (blue lines)

The linear regression relationship between PG and PD for 34 rainfall episodes, identifiable in the investigated 2014, was presented in Figure 3. Linear correlation is statistically significant ($\alpha < 0.001$).




Figure 2. Daily precipitation at 1m above the surface (PD) vs. daily precipitation above the forest canopy (PG)




Figure 3. Relationship between precipitation above the forest canopy (PG) and precipitation at 1m above the surface (PD)

Genesis of the interception phenomenon shows that the relationship between precipitation above tree crowns (PG) and precipitation at the height of 1m above the surface (PD) may be described by a following equation:

$$PD = PG + a \cdot \exp(b \cdot PG) \quad (1)$$

In the equation above (1) constants a and b are obtained in the estimation/calibration process. The results of such estimation using CurveExpert 1.4 package were presented in Figure 4. It was computed that $b = -0.01$ and $a = -3.26$. It follows that the volume of so called interception reservoir for the discussed sixty-year-old pine stand may be estimated for slightly over 3 mm (3.26).

Monthly PG and PD precipitation totals were compiled in Table 1. For the winter half year monthly PG and PD totals are similar, whereas half-year PG and PD totals are almost identical i.e. 180.0 mm each.




Figure 4. Estimated equation $PD = f(PG)$

$$PD = PG - 3.26 \cdot \exp(-0.01 \cdot PG)$$

On the other hand, for the summer half-year the differences in monthly and half year totals are apparent. Precipitation total in the summer half-year over tree crowns (PG) was 304.3 mm and rainfall at the height of 1m above the surface 209.3 mm.

Table 1. Monthly precipitation above the forest canopy (PG) and monthly precipitation at the height of 1 meter above the surface (PD)

Months	PG [mm]	PD [mm]
January	21.3	23.0
February	3.4	2.4
March	24.7	29.6
April	47.4	37.9
May	50.4	37.2
June	66.5	40.2
July	39.7	14.7
August	83.1	49.0
September	37.4	37.9
October	27.2	30.3
November	9.1	8.2
December	74.1	79.0
Jan-Dec	484.3	389.3

SUMMARY AND CONCLUSIONS

Conducted analyses allowed to formulate the following conclusions:

1. The temporary interception for the analysed sixty-year-old pine stand may reach about seven millimeters.
2. Daily difference between precipitation over the canopy and precipitation at the height of 1 above the surface may reach twelve millimeters.
3. Computed volume or so called interception reservoir was about three millimeters (3.26mm).
4. Annual value of interception for the analysed stand constitutes 19.6% of the annual precipitation total.

REFERENCES

- Bryant M.L., Bhat S., Jacobs J.M. (2005). *Measurements and modeling of throughfall variability for five forest communities in the southeastern US*. Journal of Hydrology 312, 95-108.
- Calder I.R. (2001). *Canopy processes: implications for transpiration, interception and splash induced erosion, ultimately for forest management and water resources*. Plant Ecology 153, 203-214.

- Chojnicki B. H., Urbaniak M., Danielewska A., Strzeliński P., Olejnik J. (2009). *Pomiary wymiany dwutlenku węgla oraz biomasy w ekosystemach leśnych – stacja naukowa w Tucznie*. Studia i Materiały Centrum Edukacji Przyrodniczo-Leśnej R.11, Zeszyt 2 (21), 247-256.
- Croford R.H., Richardson D.P. (2000). *Partitioning of rainfall into throughfall, stemflow and interception: Effect of forest type, ground cover and climate*. Hydrological Processes 14, 2903-2920.
- Czarnowski M.S., Olszewski J.L. (1968). *Rainfall interception by a forest canopy*. Oikos 21, Copenhagen, 48-51.
- Gash J.H.C., Loyd C.R., Lachaud G. (1995). *Estimating sparse forest rainfall interception with an analytical model*. Journal of Hydrology 170, 79-86.
- Jong de S.M., Jetten V.G. (2007). *Estimating spatial patterns of rainfall interception from remotely sensed vegetation indices and spectral mixture analysis*. [in:] International journal of geographical information science 21(5), 529-545.
- Keim R.F., Skaugset A.E., Link T.E., Iroumé A. (2004). *A stochastic model of throughfall for extreme events*. Hydrology and Earth System Sciences 8, 23-34.
- Keim R.F., Skaugset A.E., Weiler M. (2006). *Storage of water on vegetation under simulated rainfall of varying intensity*. Advances in Water Resources 29, 974-986.
- Klamerus-Iwan A. (2014). Różne spojrzenia na proces intercepcji drzew i jego determinanty. *Leśne Prace Badawcze (Forest Research Papers)*, Vol. 75 (3), 291-300.
- Olejnik J., Urbaniak M., Juszcak R., Chojnicki B., Strzeliński P., Moczko J., Roszkiewicz J., Danielewska A. (2013). *Nowatorskie i zaawansowane systemy pomiarowe*. <http://www.oic.lublin.pl/cudzechwalicie/2-3.html> [dostęp 01.03.2015].
- Osuch B., Homa A., Feliks M. (2005). *Opis pojemności zbiornika intercepcyjnego sosny zwyczajnej (Pinus silvestris)*. Czasopismo Techniczne PK.
- Pike R.G., Scherer R. (2003). *Overview of the Potential Effects of Forest Management on Low Flows in Snowmelt-dominated Hydrologic Regions*. BC Journal of Ecosystems and Management 3(1), 44-60.
- Pypker T.G., Bond B.J., Link T.E., Marks D., Unsworth M.H. (2005). *The importance of canopy structure in controlling the interception loss of rainfall: Examples from a young and an old-growth Douglas-fir forest*. Agricultural and Forest Meteorology 130, 113-129.
- Rutter A.J., Morton A.J. (1977). *A predictive model of rainfall interception in forests. III. Sensitivity of the model to stand parameters and meteorological variables*. Journal of Applied Ecology 14, 567-588.
- Suliński J. (1993). *Modelowanie bilansu wodnego w wymianie między atmosferą, drzewostanem i gruntem przy użyciu kryteriów ekologicznych*. Zeszyty Naukowe AR w Krakowie, Rozprawy habilitacyjne 179.
- Toba T., Ohta T. (2008). *Factors affecting rainfall interception determined by forest simulator and numerical model*. Hydrological Processes 22(14), 2634-2643.

Urbaniak M., Olejnik J., Miler A.T., Krysztofiak-Kaniewska A., Ziemblińska K. (2014). *Składowe bilansu wodnego w pionowym profilu dla sześćdziesięcioletniego drzewostanu sosnowego w Nadleśnictwie Tuczo*. *Infrastruktura i Ekologia Terenów Wiejskich*. Nr 2014/ II (3 (Jun 2014)), 693-704. (http://www.infraeco.pl/pl/art/a_17369.htm)

Dr inż. Marek Urbaniak

Poznań University of Life Sciences, Department of Meteorology
60-649 Poznań, ul. Piątkowska 94

Tel. 61 846-6554, E-mail: urbaniak@up.poznan.pl

Prof. dr hab. Janusz Olejnik

Poznań University of Life Sciences, Department of Meteorology
60-649 Poznań, ul. Piątkowska 94

Tel. 61 846-6550 E-mail: olejnikj@up.poznan.pl

Prof. dr hab. inż. Antoni T. Miler

Poznań University of Life Sciences, Department of Forest Engineering
60-625 Poznań, ul. Wojska Polskiego 71C

Mobile: 660-942-792, Tel. 61 848-7366, E-mail: amiler@up.poznan.pl

Dr inż. Anna Krysztofiak-Kaniewska

Poznań University of Life Sciences, Department of Forest Engineering
60-625 Poznań, ul. Wojska Polskiego 71C

Tel. 61 846-6112, E-mail: annakrysztofiak@wp.pl

Mgr inż. Klaudia Ziemblińska

Poznań University of Life Sciences, Department of Meteorology
60-649 Poznań, ul. Piątkowska 94

Tel. 61 846-6554, E-mail: klaudiaziem@wp.pl

Received: 02.01.2015

Accepted: 14.05.2015