



DOI: <https://doi.org/10.14597/infraeco.2025.001>

ASSESSMENT OF WATER REQUIREMENTS FOR ROBINIA PSEUDOACACIA L. ON RECLAIMED AREAS

Barbara JAGOSZ^{1,*}, **Wiesława KASPERSKA-WOŁOWICZ**²,
Stanisław ROLBIECKI^{3,*}, **Roman ROLBIECKI**³, **Piotr STACHOWSKI**⁴,
Wiesław PTACH⁵, **Anna FIGAS**³, **Ferenc PAL-FAM**⁶,
Daniel LIBERACKI⁴, **Ariel ŁANGOWSKI**³

ABSTRACT

Water is a crucial natural resource, influencing plant productivity and sustaining ecosystems. Black locust, known for its drought resistance and adaptability to unfavorable conditions, is frequently used in the recovery of degraded regions. During juvenile phase, black locust requires sufficient soil moisture for proper growth. The objective of the research was to assess the water needs of black locust in the second stage of growth on reclamation plantations, specifically starting three years after planting. Water requirements were evaluated for five regions in Poland between 1 June and 31 August in 1981–2010. Water requirements, quantified as crop evapotranspiration, were determined by multiplying the plant coefficient by the reference evapotranspiration. The Blaney-Criddle equation, with the plant coefficient adapted using the Żakowicz method, was applied to estimate reference evapotranspiration. It was found that the average water requirements from June to August were 370 mm. Rainfall deficits were observed in all regions, with the highest shortfalls in central Poland. Additionally, all regions demonstrated an upward trend in water requirements for this species. The results provide valuable insights for planning and implementing irrigation strategies for black locust cultivated in reclamation plantations in Poland, supporting the sustainable management of water resources in the region.

Keywords: *Black locust, irrigation, precipitation deficit, rainfall, restoration*

INTRODUCTION

The core principle of sustainable development in our natural environment is to establish an integrated framework to progressively address pre-existing imbalances and deficiencies. This involves designing systems that minimize conflicts on spatial, functional, ecological, and social levels while enhancing the overall quality of life. It also includes safeguarding the integrity of both the natural and cultural environment, optimizing the system efficiency and the effectiveness of structural development, and promoting responsible resource management, mainly concerning water resources (Koreleski, 2007).

The black locust tree is highly valued for its contributions to honey production and its ornamental qualities. It ranks among the most widely cultivated deciduous trees worldwide (Schütt *et al.*, 2006). Remarkably, this species has modest soil fertility requirements (Ellenberg and Leuschner, 2010), often thriving on abandoned or fallow lands (Başnou, 2009) and flourishing under optimal light conditions in early forest succession areas (Obidziński and Woziwoda, 2016). Its remarkability to harsh environmental conditions, such as air pollution and highly acidic soils, distinguishes the black locust (Huntley, 1990). As a result, it has been successfully used in reclamation plantings (Dilly *et al.*, 2010, Żakowicz, 2010). Recently, the cultivation of black locust for energy production has become popular, owing to its biological properties, particularly its efficient wood energy potential. The calorific value of black locust trunk wood, both in a dry and fresh state, is especially notable (Kraszkiewicz, 2016).

The black locust is an especially adaptable tree with minimal habitat requirements, often acting as a pioneer plant and displacing other woody species. Its preferred habitats typically include ruderal areas such as railway embankments, roadsides, old gravel pits, sand pits, and even rubbish dumps. The species' extensive root system is particularly remarkable, as it not only stabilizes the soil surface but also prevents erosion. In parts of Europe, particularly southern regions, the black locust's expansion has been so successful that it has begun to dominate the local tree population (Yüksek, 2012, Li *et al.*, 2018). The black locust is also widely used for establishing protective plant belts, which are vital in land reclamation efforts, providing ecological benefits and shelter for small animals (Nuța and Niculescu, 2011).

Afforestation is a fundamental approach to enhancing the function of agricultural landscapes (Pałys and Węgorek, 2005) and protecting soil and water resources (Józefaciuk *et al.*, 1995). Selecting the appropriate plant species is crucial for the success of afforestation. These species should be able to adapt easily to adverse environmental conditions, regenerate quickly after damage, exhibit low susceptibility to diseases and pests, and demonstrate rapid growth (Zajączkowski *et al.*, 2001). The ecological characteristics of the black locust closely match with these criteria (Zajączkowski *et al.*, 2001, Węgorek and Kraszkiewicz, 2005).

Black locust, known for its drought tolerance, rapid regeneration, and symbiotic relationship with nitrogen-fixing *Rhizobium* bacteria, has the potential to become the leading deciduous tree species in short rotation plantations on nutrient-poor soils. In the state of Brandenburg, Germany, situated in Central Europe with a continental climate and annual precipitation below 600 mm, the black locust has been successfully cultivated for over 250 years, primarily for wood production (Mantovani *et al.*, 2013). Black locust trees are effectively used in reclamation projects (Dilly *et al.*, 2010). *Robinia pseudoacacia* L. is commonly employed for the biological reclamation of post-industrial wastelands and degraded lands (Rahmonov, 2009). Notably, black locust thrives even in reclaimed post-mining areas characterized by spring-to-summer drought and poor soil water availability (Mantovani *et al.*, 2014a).

Despite its modest water needs, young black locust trees need adequate soil moisture until they establish a symbiotic relationship with *Rhizobium* bacteria (Rédei *et al.*, 2008). In Poland, *Robinia pseudoacacia* L. is widely used for reclaiming a significant portion of degraded lands. Its applications include forest reclamation on the slopes of external dumps from open-cast mines and nutrient-poor, loose sands. Black locust plays a crucial role in

stabilizing substrates and improving soil nutrient content. Areas reclaimed with black locust are effectively protected against wind and water erosion. The leaves of the black locust provide a primary source of organic matter, fostering humus layer development and enriching the soil with essential macro- and micro-elements (Rahmonov and Parusel, 2012). Black locust forests enhance nutrient-deficient substrates with nutrients and organic colloids, significantly influencing the progress of ecosystem processes, thereby contributing to habitat diversity (Read *et al.*, 2002, Tamminen *et al.*, 2004).

The extensive use of black locust in reclamation plantings has significant potential for bioenergy production (Cafourek *et al.*, 2016). Ensuring the survival of plantings in reclaimed areas is a fundamental strategy during the initial phase of reclamation. Successful development of reclamation plantings requires two key elements: selecting the appropriate species and planting high-quality seedlings (Klimek *et al.*, 2008, 2009, 2011, 2013), as well as providing adequate soil moisture through irrigation (Żakowicz, 2010). While mature black locust trees can readily adapt to dry soil conditions, trees in the juvenile phase require optimal soil moisture levels. This can be achieved through irrigation techniques that compensate for precipitation deficits (Żakowicz, 2010, Mantovani *et al.*, 2014 a, b).

The primary aim of this study was to estimate the water requirements of black locust trees during the second stage of growth on reclamation plantations, specifically in the period beginning three years after reclamation. The research focused on the three-month period of increased water needs for black locust, occurring from June to August. This study builds upon previous research by Rolbiecki *et al.* (2019), who evaluated the water requirements of young black locust seedlings during the first phase of development on reclamation plantations, from planting until three years after planting.

MATERIAL AND METHOD

This study calculated the water requirements of black locust (*Robinia pseudoacacia* L.) over three years following planting in a reclaimed area, assuming the plant's water needs equivalent to potential crop evapotranspiration under sufficient soil moisture conditions. The water needs of young black locust trees were assessed using the crop coefficient method, which involves multiplying the crop coefficient by the reference evapotranspiration. The calculations were carried out using Equation (1):

$$ET_p = ETo \times kc \quad (1)$$

where:

ET_p – represents potential crop evapotranspiration (mm)

ETo – denotes reference evapotranspiration (mm)

kc – is the crop coefficient, defined as the ratio of evapotranspiration measured under adequate soil moisture conditions to reference evapotranspiration (Łabędzki, 2006).

The Blaney-Criddle formula, adjusted to Polish conditions by Żakowicz (2010), was used to calculate the reference evapotranspiration. Equation (2) was applied for the calculations:

$$ET_o = n \times [p \times (0.437 \times t + 7.6) - 1.5] \quad (2)$$

where:

ET_o – represents reference evapotranspiration (mm)

n – is the number of days in the month

p – refers to evaporation coefficients by Doorenbos and Pruitt (1977) for specific months and latitude, as determined from tables, t – indicates the monthly mean air temperature ($^{\circ}\text{C}$).

For young black locust trees growing in reclaimed areas, starting three years after reclamation, crop coefficients were applied according to the guidelines established by Żakowicz (2010).

The calculations were conducted for the period from 1 June to 31 August, which corresponds to the time when plants in Poland have the highest water requirements during the growing season. The water requirements of black locust trees were estimated based on data collected over a thirty-year span, from 1981 to 2010, from five meteorological stations located in Olsztyn, Bydgoszcz, Warszawa, Wrocław, and Kraków. These stations represent various agro-climatic regions of Poland, including the north-east region (N-E), central-north-west region (C-N-E), central-east region (C-E), south-west region (S-W), and south-east region (S-E), respectively (Figure 1).




Figure 1. Geographical position of the investigated regions in Poland

The rainfall deficit for young black locust trees was calculated by subtracting the cumulative precipitation from the total potential crop evapotranspiration. To evaluate the rainfall deficit under different probabilities, including normal years ($N_{50\%}$), medium dry years ($N_{25\%}$), and very dry years ($N_{10\%}$), Ostromęcki's method was applied (Żakowicz *et al.*, 2009) using Equation (3):

$$Np\% = Ap\% \times ETp - Bp\% \times P \quad (3)$$

where:

$Np\%$ – represents the precipitation deficit at the occurrence probability $p\%$ (mm per period)

$Ap\%$ and $Bp\%$ – are numerical factors that describe the variability of evapotranspiration and precipitation for a specific meteorological station

ETp – signifies the multi-year average evapotranspiration during the study period (mm per period)

P – denotes the multi-year average precipitation for the same period (mm per period).

RESULT AND DISCUSSION

The variation in water requirements of young black locust trees, as indicated by the coefficient of variation computed over the thirty-year period between 1981 and 2010 for the months of peak water demand—June, July, and August—showed relatively low values, ranging from 3.7% in the central–east region to 4.1% in the south–west region of Poland (Table 2).

Table 2. Characteristics of black locust water requirements from June to August, 1981–2010

Characteristic	Region of Poland	Water needs
Minimum (mm)	north–east	340.0
	central–north–west	358.8
	central–east	354.3
	south–west	330.9
	south–east	327.9
Maximum (mm)	north–east	393.2
	central–north–west	417.8
	central–east	415.6
	south–west	391.2
	south–east	388.2
Median (mm)	north–east	365.9
	central–north–west	385.9
	central–east	384.1
	south–west	357.7
	south–east	358.7
Standard deviation (mm)	north–east	14.3
	central–north–west	15.2
	central–east	14.2
	south–west	14.6
	south–east	13.5
Variation coefficient (%)	north–east	3.9
	central–north–west	3.9
	central–east	3.7
	south–west	4.1
	south–east	3.8

During the analyzed period from 1981 to 2010, a noticeable trend of increasing water requirements for black locust trees from 1 June 31 to August was observed across all regions across Poland (Figures 2 a–f). The trend lines depicting the variability of black locust water requirements (R^2) demonstrated a significant upward trajectory throughout the country ($R^2 = 0.2556$) (Figure 2 f). The lowest R^2 value for this trend was observed in the central–north–west

region of the country ($R^2 = 0.0907$) (Figure 2 b). Conversely, in the north-east region ($R^2 = 0.2525$) (Figure 2 a), south-west region ($R^2 = 0.2674$) (Figure 2 d), central-east region ($R^2 = 0.2734$) (Figure 2 c), and south-east region ($R^2 = 0.3447$) (Figure 2 e) of Poland, the trend of increasing water requirements for black locust trees was notably more pronounced. The most significant increase in water requirements was observed in the south-east region of Poland, with an increase of 9.1 mm per decade (Figure 2 e). In contrast, the smallest rise in water needs occurred in the central-north-west region, with an increase of 5.3 mm per decade (Figure 2 b). On average, across Poland, from 1981 to 2010, during the period from 1 June to 31 August, the water requirements of black locust increased by an average of 8 mm per decade (Figure 7).





Figure 2. The time trend analysis of black locust water requirements in the particular regions of Poland. The solid line corresponds to the trend line

The most substantial water needs for black locust trees, as considered in the current study, were observed in the central regions of Poland. During the three-decade period from June to August, the highest water needs were recorded in the central-north-west (386 mm) and central-east (384 mm) regions of the country (Figure 3). In contrast, the lowest water needs were calculated in the south-west (359 mm) and south-east (357 mm) regions of Poland. Several studies affirm the suitability of central Poland for agricultural (Stachowski and Markiewicz 2011, Żarski *et al.*, 2013), horticultural (Żarski *et al.*, 2013, Rolbiecki *et al.*, 2002 a, b), and forestry crops (Rolbiecki *et al.*, 2018), despite these areas experiencing the highest rainfall deficits and having

soil with low water retention. These studies concurrently emphasize the necessity of irrigation for productive land use in these regions.

According to Żakowicz *et al.* (2009), precipitation deficit calculations were made based on the probability of occurrence for normal years ($N_{50\%}$, i.e., once every two years), medium dry years ($N_{25\%}$, i.e., once every four years), and very dry years ($N_{10\%}$, i.e., once every ten years). In these three categories of years, the plants' water demand is met by 50%, 75%, and 90%, respectively. In the current study, from 1 June to 31 August, the largest rainfall deficit for normal years ($N_{50\%}$) was observed in the central–north–west region of Poland (197 mm) and the central–east region (173 mm), while the lowest was recorded in the south–east region (105 mm) of the country (Table 3). On the other hand, the rainfall deficit in medium dry years ($N_{25\%}$) and very dry years ($N_{10\%}$) was significantly greater in the north–east (277 mm and 364 mm, respectively), central–north–west (287 mm and 356 mm, respectively) and central–east regions (281 mm and 402 mm, respectively), compared to the south–west (213 mm and 301 mm, respectively) and south–west regions (191 mm and 242 mm, respectively) of the country.



Regions and representative meteorological stations

Figure 3. Black locust water needs for the June–August period in various regions of Poland: N–E, north–east; C–N–E, central–north–west; C–E, central–east; S–W, south–west and S–E, south–east region, PL, Poland

Table 3. Rainfall deficit (mm) calculated for black locust trees in various regions of Poland

Probability of rainfall deficit occurrence	Region of Poland				
	North–East	Central–North–West	Central–East	South–West	South–East
Normal years	150	197	173	141	105
Medium dry years	277	287	281	213	191
Very dry years	364	356	402	301	242

On average, the calculated water needs for black locust trees in Poland from 1 June to 31 August were 370 mm, as depicted in Figure 3. The natural habitat of black locust trees originally spans a humid to sub-humid climatic region in North America, characterized by an annual precipitation ranging from 1020 mm to 1830 mm. Remarkably, black locust trees exhibit a high

degree of drought tolerance and can thrive across a wide range of climatic and soil conditions (Mantovani *et al.*, 2013, 2015).

Globally, this tree species plays a significant role in enhancing soil fertility and positively influencing the chemical properties of the soil (Bolat *et al.*, 2016), contributing significantly to the reclamation of degraded soils (Mantovani *et al.*, 2015, Quinkenstein *et al.*, 2011, Nicolescu *et al.*, 2018). In central Europe, which has a continental climate, particularly in marginal areas like post-mining lands with limited water resources and challenging soil conditions, black locust cultivation for biomass production has been successful (Mantovani *et al.*, 2013). It's noteworthy that in these post-mining regions, where pre-dawn water potential as low as -0.5 MPa was observed during summer droughts, stem growth was severely limited. Nevertheless, black locust plants managed to survive these adverse weather and soil conditions (Mantovani *et al.*, 2015).

The results presented in this study indicate a growing trend in the water needs of black locust trees from 1 June to 31 August across Poland. Rapid climate changes, including rising temperature, may further increase the water needs of these plants in the near future. In light of this, plants developed in reclamation areas will be particularly vulnerable to adverse climate changes. Consequently, adaptation measures, including irrigation, become crucial and are among the most significant agro-technical practices that support plant development when rainfall is insufficient (Rolbiecki *et al.*, 2018, Kuchar and Iwański, 2013, Kuchar *et al.*, 2015, 2017). According to Ptach *et al.* (2018), irrigation is widely recognized as a critical melioration technique that facilitates the growth of trees and shrubs in various planting settings, including forests and tree nurseries. Numerous experiments conducted in central Poland have demonstrated the positive impact of irrigation on the seedling development of various tree species, including Scots pine (Klimek *et al.*, 2008), white birch (Klimek *et al.*, 2009), European larch (Klimek *et al.*, 2011), little leaf linden (Klimek *et al.*, 2013), and paulownia (Ptach *et al.*, 2018).

Earlier studies conducted by Żakowicz (2010) and Żakowicz and Hewelke (2012) have already established the effectiveness of micro-sprinkler irrigation for black locust grown in reclaimed areas in central Poland. These studies observed that irrigation using water from sewage treatment plants or rainwater positively influenced the height, trunk area, and crown area of black locust plants. Additionally, Mantovani *et al.* (2013, 2014 a, b) repeatedly noted the positive effects of drip irrigation on the growth and development of two-year-old black locust trees.

CONCLUSION

The study presented here indicates that, in the reclaimed areas in Poland, the average water requirements for young black locust trees during the three summer months (June, July, August) starting three years after planting, amounted to 370 mm. Rainfall deficits were calculated across Poland, with the most significant deficits observed in the central–north–west and central–east regions. Furthermore, an upward trend in water needs is evident. Black locust trees, known for their drought tolerance, are commonly used in the ecological restoration of degraded areas. However, during the early stages of growth, these trees require optimal soil moisture levels to ensure proper development, which is a crucial consideration for reclamation projects on degraded lands. The water requirement values obtained from this study can be valuable in developing a program for supplementary irrigation treatments in

the reclamation plantings of young black locust trees, aimed at optimizing their growth in Poland. This strategy will facilitate the efficient management of water resources in the studied area. Future research should focus on evaluating the impact of irrigation on black locust trees during periods of water deficit.

REFERENCES

1. Başnou, C. (2009). *Robinia pseudoacacia*, black locust (Fabaceae, Magnoliophyta). [In:] Drake J.A. ed. *Handbook of alien species in Europe*. Dordrecht, Netherlands: Springer.
2. Bolat, I., Kara, O., Sensoy, H., Yuksel, K. (2016). *Influences of black locust (Robinia pseudoacacia L.) afforestation on soil microbial biomass and activity*. iForest 9: 171-177. DOI: 10.3832/for1410-007
3. Cafourek, J., Gaff, M., Gašparík, M., Slávik, M., Macků, J. (2016). *Properties and use of biomass from reclaimed land in the North Bohemian Basin*. Wood Research 61(5): 777-790.
4. Dilly, O., Nii-Annang, S., Schrautzer, J., Breuer, V., Pfeiffer, E.M., Gerwin, W., Schaaf, W., Freese, D., Veste, M., Huttel, R.F. (2010). *Ecosystem manipulation and restoration on the basis of long-term conceptions*. [In:] Müller F., Baessler C., Schubert H., Klotz S. eds. *Long-Term Ecological Research Between Theory and Application*. Amsterdam, Netherlands: Springer Science+Business Media: 411-428.
5. Doorenbos, J., Pruitt, W.O. (1977). *Guidelines for predicting crop water requirements*. FAO Irrigation and Drainage Paper 24: 176.
6. Ellenberg, H., Leuschner, C. (2010). *Vegetation Mitteleuropas mit den Alpen in ökologischer, dynamischer und historischer Sicht*. Stuttgart, Germany: H. Eugen Ulmer.
7. Huntley, J.C. (1990). *Robinia pseudoacacia L. Black locust*. [In:] Burns R.M., Honkala B.H. eds. *Silvics of North America, Vol. 2. Hardwoods*, Agricultural Handbook. U.S. Washington, US, Department of Agriculture, Forest Service, 654: 755-761.
8. Józefaciuk, C., Tałałaj, Z., Węgorzek, T. (1995). *Ochronno-produkcyjna funkcja zadrzewień pasowych na przykładzie obiektu Linów*. Pamiętnik Puławski 106: 173-182.
9. Klimek, A., Rolbiecki, S., Rolbiecki, R., Hilszczańska, D., Malczyk, P. (2008). *Impact of chosen bare root nursery practices in Scots pine seedling quality and soil mites (Acari)*. Polish Journal of Environmental Studies 17(2): 247-255.
10. Klimek, A., Rolbiecki, S., Rolbiecki, R., Malczyk, P. (2009). *Impact of chosen bare root nursery practices on white birch seedling quality and soil mites (Acari)*. Polish Journal of Environmental Studies 18(6): 1013-1020.
11. Klimek, A., Rolbiecki, S., Rolbiecki, R., Hilszczańska, D., Malczyk, P. (2011). *Effects of organic fertilization and mulching under micro-sprinkler irrigation on growth and mycorrhizal colonization of European larch seedlings, and occurrence of soil mites*. Polish Journal of Environmental Studies 5(20): 1211-1219.
12. Klimek, A., Rolbiecki, S., Rolbiecki, R., Długosz, J., Musiał, M. (2013). *The use of compost from sewage sludge and forest ectohumus for enrichment of soils in the nursery cultivation of littleleaf linden (Tilia cordata Mill.)*. Annual Set The Environment Protection 15: 2811-2828.

13. Koreleski, K. (2007). *Koncepcja rozwoju zrównoważonego w unijnej polityce kształtowania obszarów wiejskich*. Infrastructura and Ecology of Rural Areas 1: 19-26.
14. Kraszkiewicz, A. (2016). Chemical composition and selected energy properties of black locust bark (*Robinia pseudoacacia* L.). *Agricultural Engineering* 20(2): 117-124. DOI: 10.1515/agriceng-2016-0033
15. Kuchar, L., Iwański, S. (2013). *Rainfall evaluation for crop production until 2050-2060 and selected climate change scenarios for North Central Poland*. Infrastructura and Ecology of Rural Areas 2(I): 187-200.
16. Kuchar, L., Iwański, S., Diakowska, E., Gąsiorek, E. (2015). *Simulation of hydrothermal conditions for crop production purpose until 2050-2060 and selected climate change scenarios for North Central Poland*. Infrastructura and Ecology of Rural Areas II(1): 319-334.
17. Kuchar, L., Iwański, S., Diakowska, E., Gąsiorek, E. (2017). *Assessment of meteorological drought in 2015 for North Central part of Poland using hydrothermal coefficient (HTC) in the context of climate change*. Infrastructura and Ecology of Rural Areas I(2): 257-273.
18. Li, G., Zhang, X., Huang, J., Wen, Z., Du, S. (2018). *Afforestation and climatic niche dynamics of black locust (Robinia pseudoacacia)*. *Forest Ecology and Management* 407: 184-190. DOI: 10.1016/j.foreco.2017.10.019
19. Łabędzki, L. (2006). *Susze rolnicze. Zarys problematyki oraz metody monitorowania i klasyfikacji*. Woda. Środowisko. Obszary Wiejskie. Rozprawy Naukowe i Monografie 17: 1-107.
20. Mantovani, D., Veste, M., Boldt-Burisch, K., Fritsch, S.; Freese, D. (2013). *Black locust (Robinia pseudoacacia L.) root growth response to different irrigation regimes*. *Verhandlungen der Gesellschaft für Ökologie*. 43: 333.
21. Mantovani, D., Veste, M., Freese, D. (2014 a). *Black locust (Robinia pseudoacacia L.) adaptability and plasticity to drought*. 2nd European Agroforestry Conference: Integrating Science and Policy to Promote Agroforestry in Practice, Cottbus, Germany, 4-6 June, Book of Abstracts, pp. 264-265.
22. Mantovani, D., Veste, M., Freese, D. (2014 b). *Black locust (Robinia pseudoacacia L.) ecophysiological and morphological adaptations to drought and their consequence on biomass production and water-use efficiency*. *New Zealand Journal of Forestry Science* 44: 29. DOI: 10.1186/s40490-014-0029-0
23. Mantovani, D., Veste, M., Böhm, C., Vignudelli, M., Freese, D. (2015). *Spatial and temporal variation of drought impact on black locust (Robinia pseudoacacia L.) water status and growth*. *iForest* 8: 743-747. DOI: 10.3832/ifor1299-008
24. Nicolescu, V.N., Hernea, C., Bakti, B., Keserü, Z., Antal, B., Rédei, K. (2018). *Black locust (Robinia pseudoacacia L.) as a multi-purpose tree species in Hungary and Romania: a review*. *Journal of Forestry Research* 29: 1449-1463. DOI: 10.1007/s11676-018-0626-5
25. Nuța, S.I., Niculescu, M. (2011). *The influence of forest belts on tobacco crops in hydro-ameliorative Sadova-Corabia system*. *Analele Universității din Craiova, seria Agricultura-Montanologie-Cadastru, Craiova* XLI(2): 210-214.
26. Obidziński, A., Woziwoda, B. (2016). *Robinia akacjowa (Robinia pseudoacacia) [Black locust (Robinia pseudoacacia)]*. [In:] Obidziński A., Kołaczowska E., Otręba A. eds. *Metody zwalczania obcych*

- gatunków roślin występujących na terenie Puszczy Kampinoskiej. Izabelin–Kraków, Poland: BioDar: 106-121.
27. Pałys, S., Węgorek, T. (1988). *Melioracje przeciwerozyjne a ochrona środowiska*. Wiadomości Melioracyjne i Łąkarskie 5-6: 135-137.
 28. Ptach, W., Łangowski, A., Rolbiecki, R., Rolbiecki, S., Jagosz, B., Grybauskiene, V., Kokoszewski, M. (2018). *The influence of irrigation on the growth of paulownia trees at the first year of cultivation in a light soil*. Proceedings of the 8th International Scientific Conference Rural Development 2017, Bioeconomy Challenges, Lithuania, 23-24 November, 2017; Raupelienė, A. ed. Aleksandras Stulginskis University: 763-768.
 29. Quinkenstein, A., Boehm, C., Matos, E., Freese, D., Huettl, R.F. (2011). *Assessing the carbon sequestration in short rotation coppices of Robinia pseudoacacia L. on marginal sites in Northeast Germany*. [In:] Kumar B.M., Nair P.K.K. eds. Carbon sequestration potential of agroforestry systems: opportunities and challenges. New York, US: Springer: 201-216.
 30. Rahmonov, O. (2009). *The chemical composition of plant litter of black locust (Robinia pseudoacacia L.) and its ecological role in sandy ecosystems*. Acta Ecologica Sinica 29: 237-243. DOI: 10.1016/j.chnaes.2009.08.006
 31. Rahmonov, O., Parusel, T. (2012). *Wpływ opadu roślinnego robinii akacjowej Robinia pseudoacacia L. na proces rozwoju gleby na obszarach zdegradowanych*. Studia i Materiały CEPL w Rogowie, R14(33/4): 81-92.
 32. Read, J., Ferris, J.M., Jaffre T. (2002). *Foliar mineral content of Nothofagus species on ultramafic soils in New Caledonia and non-ultramafic soils in Papua New Guinea*. Australian Journal of Botany 50(5): 607-617. DOI.org/10.1071/BT01091
 33. Rédei, K., Osváth-Bujtás, Z., Veperdi, I. (2008). *Black locust (Robinia pseudoacacia L.) improvement in Hungary: a review*. Acta Silvatica et Lignaria Hungarica 4: 127-132. DOI: 10.37045/aslh-2008-0011
 34. Rolbiecki, S., Rolbiecki, R., Rzekanowski, C. (2002 a). *Response of black currant (Ribes nigrum L.) cv. 'Titania' to micro-irrigation under loose sandy soil conditions*. Acta Horticulturae 585(2): 649-652. DOI: 10.17660/ActaHortic.2002.585.107
 35. Rolbiecki, S., Rolbiecki, R., Rzekanowski, C. (2002 b). *Effect of micro-irrigation on the growth and yield of raspberry (Rubus idaeus L.) cv. 'Polana' grown in very light soil*. Acta Horticulturae 585(2): 653-657. DOI: 10.17660/ActaHortic.2002.585.108
 36. Rolbiecki, S., Kokoszewski, M., Grybauskiene, V., Rolbiecki, R., Jagosz, B., Ptach, W., Łangowski A. (2018). *Effect of expected climate changes on the water needs of forest nursery in the region of central Poland*. Proceedings of the 8th International Scientific Conference Rural Development 2017, Bioeconomy Challenges, Lithuania, 23-24 November, 2017; Raupelienė, A. ed. Aleksandras Stulginskis University: 786-792.
 37. Rolbiecki, S., Kasperska-Wołowicz, W., Ptach, W., Jagosz, B., Figas, A., Rolbiecki, R., Stachowski, P., Grybauskiene, V., Chmura, K., Dobosz, K. (2019). *Water Needs of Black Locust Robinia pseudacacia L. in the First Three Years of Growing in the Reclamation Plantings in Poland*. [In:] Infrastructure and Environment. Berlin/Heidelberg, Germany, Springer International Publishing: 234-242.

38. Schütt, P., Weisgerber, H., Schuck, H.J., Lang, U.M., Stimm, B., Roloff, A. (2006). *Enzyklopädie der Laubbäume: Die große Enzyklopädie*. Hamburg, Germany: Nikol Verlag Barkhausenweg.
39. Stachowski, P., Markiewicz, J. (2011). *The need of irrigation in central Poland on the example of Kutno county*. Annual Set The Environment Protection 13: 1453-1472.
40. Tamminen, P., Starr, M., Kubin, E. (2004). *Element concentrations in boreal, coniferous forest humus layers in relation to moss chemistry and soil factors*. Plant and Soil 259: 51-58.
41. Węgorzek, T., Kraszkiewicz, A. (2005). *Dynamika wzrostu robinii akacjowej (Robinia pseudacacia L.) w zadrzewieniu śródpolnym na glebach lessowych*. Acta Agrophysica 5(1): 211-218.
42. Yükses, T. (2012). *The restoration effects of black locust (Robinia pseudoacacia L.) plantation on surface soil properties and carbon sequestration on lower hillslopes in the semi-humid region of Coruh Drainage Basin in Turkey*. Catena 90: 18-25. DOI.org/10.1016/j.catena.2011.10.001
43. Zajączkowski, K., Tałaaj, Z., Węgorzek, T., Zajączkowska, B. (2001). *Dobór drzew i krzewów do zadrzewień na obszarach wiejskich*. Warszawa, Poland: IBL.
44. Żakowicz, S. (2010). *Podstawy technologii nawadniania rekultywowanych składowisk odpadów komunalnych*. Rozprawy Naukowe i Monografie. Warszawa, Poland: SGGW.
45. Żakowicz, S., Hewelke, P., Gnatowski, T. (2009). *Podstawy infrastruktury technicznej w przestrzeni produkcyjnej*. Warszawa, Poland: SGGW.
46. Żakowicz, S., Hewelke, P. (2012). *Technologia nawadniania roślin na rekultywowanych składowiskach odpadów komunalnych*. Warszawa, Poland: SGGW.
47. Żarski, J., Dudek, S., Kuśmierk-Tomaszewska, R., Rolbiecki, R., Rolbiecki, S. (2013). *Forecasting effects of plants irrigation based on selected meteorological and agricultural drought indices*. Annual Set The Environment Protection 15: 2185-2203.

Corresponding author: Barbara Jagosz

ORCID: 0000-0003-2701-8246

e-mail: Barbara.Jagosz@urk.edu.pl

University of Agriculture in Krakow

Faculty of Biotechnology and Horticulture

Department of Plant Biology and Biotechnology

Mickiewicza 21, 31-120 Kraków, Poland

Wiesława Kasperska-Wołowicz

ORCID: 0000-0001-8265-8786

e-mail: w.kasperska-wolowicz@itp.edu.pl

Institute of Technology and Life Sciences

Kuyavian-Pomeranian Research Centre

Glinki 60, 85-174 Bydgoszcz, Poland

Corresponding author: Stanisław Rolbiecki

ORCID: 0000-0002-1433-2212

e-mail: rolbs@pbs.edu.pl

Bydgoszcz University of Science and Technology

Faculty of Agriculture and Biotechnology

Department of Agrometeorology, Plant Irrigation and Horticulture

Sylwestra Kaliskiego 7, 85-796 Bydgoszcz, Poland

Roman Rolbiecki

ORCID: 0000-0001-6230-4227

e-mail: rolbr@pbs.edu.pl

Bydgoszcz University of Science and Technology

Faculty of Agriculture and Biotechnology

Department of Agrometeorology, Plant Irrigation and Horticulture

Sylwestra Kaliskiego 7, 85-796 Bydgoszcz, Poland

Piotr Stachowski

ORCID: 0000-0002-1367-0551

e-mail: piotr.stachowski@up.poznan.pl

Poznań University of Life Sciences

Faculty of Environmental Engineering and Mechanical Engineering

Department of Land Improvement, Environmental Development and Spatial

Management

Piątkowska 94, 60-649 Poznań, Poland

Wiesław Ptach

ORCID: 0000-0002-7007-0931

e-mail: wieslaw_ptach@sggw.edu.pl

Warsaw University of Life Sciences

Institute of Environmental Engineering

Department of Remote Sensing and Environmental Research

Nowoursynowska 159, 02-776 Warszawa, Poland

Anna Figas

ORCID: 0000-0002-9021-8702

e-mail: figasanna@pbs.edu.pl

Bydgoszcz University of Science and Technology

Faculty of Agriculture and Biotechnology

Department of Agricultural Biotechnology

Sylwestra Kaliskiego 7, 85-796 Bydgoszcz, Poland

Ferenc Pal-Fam

ORCID: 0000-0002-1545-7885

e-mail: Pal-Fam.Ferenc.Istvan@uni-mate.hu

Hungarian University of Agriculture and Life Sciences

Institute of Plant Production

Kaposvár Campus, H 7400 Kaposvár, Hungary

Daniel Liberacki

ORCID: 0000-0002-4582-4535

e-mail: daniel.liberacki@up.poznan.pl

Poznań University of Life Sciences

Faculty of Environmental Engineering and Mechanical Engineering

Department of Land Improvement, Environmental Development and Spatial

Management

Piątkowska 94, 60-649 Poznań, Poland

Ariel Łangowski

ORCID: 0000-0002-3459-8990

e-mail: arilan000@pbs.edu.pl

Bydgoszcz University of Science and Technology

Faculty of Agriculture and Biotechnology

Department of Agrometeorology, Plant Irrigation and Horticulture

Sylwestra Kaliskiego 7, 85-796 Bydgoszcz, Poland

Received: 02.12.2025

Revised: 02.02.2025

Accepted: 05.02.2025